




Phaeochromocytomas and Paragangliomas

NET Patient Foundation and AMEND





Phaeochromocytomas & Paragangliomas

(incl MEN and SDHx)

Phaeochromocytomas, often known as 'phaeos' ('fee-ohs'), are rare tumours of the adrenal gland arising from the inner section of the gland called the medulla. Similar tumours can arise from sites other than the adrenal gland and these are called extra-adrenal paragangliomas.

They can affect people of all ages, including children. They are classified as a type of neuroendocrine tumour (NET) as they secrete certain peptides and hormones. The term phaeochromocytoma is derived from the greek words 'phaeos' meaning dusky, 'chromo' meaning colour and 'cytoma' meaning tumour. It refers to the brownish colour the tumour cells develop when the tissue is treated with special chemicals to view them under a microscope.

The body has two walnut-sized adrenal glands, one above each of the kidneys ('adrenal' means next to the kidney). Even though the glands are small they are important as they produce several essential hormones. The medulla (the area of the adrenal gland from which phaeos arise) produces a number of hormones called catecholamines, mainly adrenaline and noradrenaline. These hormones help the body deal with maintaining blood pressure, deal with sudden stress or threats. Phaeos may produce excessive amounts of these hormones, causing a variety of symptoms that can affect the whole body.

What is known about these tumours?

- This type of tumour is rare occurring in 1 per 100 000 people per year
- The majority (approximately 90% or 9 out of 10) are benign tumours (i.e. not cancer)
- Over 80% (8 in 10 cases) arise in the adrenal glands
- The most common age for diagnosis is between the ages of 30-60 years of age, although 10% arise in children
- There is no sex difference or racial disposition.
- Tumours occurring in children are more likely to be related to an inherited condition than those occurring in adults.



Is there a genetic link?

- The majority of these tumours are not linked to a genetic condition and are therefore known as 'sporadic' tumours.
- Approximately 25% (1 in 4 cases) are related to inheriting a gene change which leads to an increased risk of developing phaeochromocytoma.
- The inherited forms can occur in syndromes, including: multiple endocrine neoplasia 2 (MEN-2), neurofibromatosis and von-hippel lindau syndrome.
- There are families who develop phaeos with paragangliomas due to a very rare change in the succinate dehydrogenase (SDH) gene (see 'Inherited Paraganglioma Syndromes').
- Multiple Endocrine Neoplasia type 2 (MEN2): approximately 50% (1 in 2) of patients with a gene change that causes MEN2 will develop phaeos, often in both adrenal glands though not necessarily at the same time. They are almost always benign (not cancer) in MEN and would be treated in the same way as a sporadic phaeo. In addition to phaeos, patients with a MEN2 gene change almost always develop a rare form of thyroid cancer (medullary thyroid cancer) and less commonly, benign tumours of the parathyroid glands in the neck which may cause raised levels of calcium in the blood. Visit the Association for Multiple Endocrine Neoplasia Disorders (AMEND) website at www.amend.org.uk for more information on MEN2.

What are the symptoms of phaeochromocytomas?

Phaeos can cause a wide variety of symptoms which are mainly related to the release by the tumour of excessive amounts of hormone, mainly adrenaline and noradrenaline. Symptoms can occur intermittently often in episodes lasting only 15 minutes and may include some or all of the following:

- Headaches
- Dizziness
- Facial paleness (pallor)
- Excessive sweating
- Racing heart rate (palpitations)
- Panic attacks/ sense of doom
- Anxiety
- Weight loss
- Heat intolerance
- High (and rarely low) blood pressure (sustained or episodic)
- Nausea (with or without vomiting)
- Breathlessness
- Depression
- Lethargy

Many of the symptoms listed above are related to the high blood pressure (hypertension) that can be caused due to this condition. It is thought one case of hypertension in every 1000-2000 patients is due to phaeochromocytoma. Hypertension in patients with undiagnosed phaeos can be very difficult to control; however, once a diagnosis is made, special drugs are available to help to control it.



Phaeos in children

In children most phaeos occur in just one adrenal gland, however, in 20% (1 in 5) of cases they can affect both glands.

Approximately 1 in 3 young patients have tumours outside the adrenal gland with or without a tumour inside the gland.

The initial symptoms in children are similar to those of adults, including headache, fast heart rate and sweating. Their blood pressure is often raised. Children require the same investigations as adults to determine the underlying cause and make a diagnosis.

How are phaeos diagnosed?

Unless a patient is known to have a higher risk of developing tumours due to an inherited condition such as MEN2, phaeos are notoriously hard to diagnose as the symptoms may be very varied and often occur in episodes or sudden attacks lasting less than 15 minutes. If a phaeo is suspected a number of various tests may be recommended by your doctor. These may include:

24 hour urine test (catecholamines): This test measures the amount of catecholamines that are excreted in the urine over a 24 hour period. In phaeos this level will be much higher than normal. The collection bottles contain acid as a preservative and should not be refrigerated. This test is good in helping diagnose phaeos but is not 100% accurate and so multiple collections may be required as well as other tests listed below.

24 hour urine test (metanephrines): This test is similar to the one above but measures the level of metanephrines in the urine excreted over 24 hours. Metanephrines are breakdown products of adrenaline and noradrenaline. The urine collection bottles do not contain acid and should not be refrigerated.

Plasma Metanephrine Test: Increasingly, plasma (blood) metanephrine and normetanephrine tests are being used to identify the presence of a phaeo. This test should be performed after the patient has been lying quietly for about 30 minutes to avoid a false positive result.

CT scan: a computer tomography scan provides a three dimensional picture of the inside of the body. It can be used to determine the position and size of tumours.

MRI scan: This magnetic resonance imaging (MRI) scan can help reveal where the tumours are positioned. It uses magnetism rather than x-rays to take pictures of the inside of the body.

MIBG scan: a specialised scan performed at the hospital's nuclear medicine department. MIBG (Meta iodo benzyl guanidine) is a chemical that is readily picked up by many phaeos. When the MIBG is combined with a mildly radioactive agent and injected via a vein in the arm, it sticks to the tumour cells which light up on the screen as radioactive hotspots.

PET imaging: a PET (positron emission tomography) scan is another nuclear medicine imaging technique similar to MIBG but which uses different agents that can either bind or be taken up by the tumour. Currently, MIBG remains the nuclear medicine imaging standard, however, as more research is done this may change.



What is the treatment?

Treatment depends on many factors such as the location of the tumour and whether it has spread.

Most phaeos are benign, especially those occurring in patients with MEN-2. This means they are not cancerous and do not spread. If it is possible to remove the tumour completely, this may be the only treatment required. Surgery usually involves the removal of the entire adrenal gland and is called an adrenalectomy. If your phaeo is believed to be cancer and it has spread locally, then the surgeon will remove the tissues immediately around the adrenal gland and the nearby lymph nodes. If one adrenal gland is removed, then other gland will produce enough hormones such that you will not need to take medication. However, if both adrenal glands are removed, then you will need life-long medication in the form of steroids to replace the hormones that the adrenal glands would normally make.

Due to the potential for sudden dangerous spikes in blood pressure caused when phaeos are handled during surgery, patients are given special blood pressure medications before surgery. This will sometimes be done as an inpatient. Also occasionally radionuclide therapy may be used to control symptoms.

Surgery

This depends on the size of the tumour, and whether one or both adrenal glands are being removed. Most tumours can be removed laparoscopically, which means using minimally invasive keyhole surgery performed via a series of small incisions in the abdomen. Some larger tumours may be removed either through a single incision in the abdomen or through the back removing a lower rib.

Some expressions you may hear are:

- Right hand (RH) adrenalectomy: removal of the right side adrenal gland only.
- Left hand (LH) adrenalectomy: removal of the left side adrenal gland only.
- Bilateral adrenalectomy: the removal of both adrenal glands at the same time.

Phaeos can cause very unstable blood pressure during and after surgery. To help stabilize this you will be given medication (anti-hypertensive drugs) called alpha- and sometimes also beta- blockers for up to 10 days before surgery. This practise of controlling the blood pressure is done even for patients with little or no symptoms in order to minimize any risk during surgery. Following surgery the blood pressure should return to normal.

Targeted radionuclide therapy

If surgery is not an option, targeted radioactive therapy using I^{131} -MIBG is sometimes an option. As 90% of these tumours are positive to an MIBG scan, they are often suitable for this type of targeted therapy. You may have already had an MIBG scan during the tests requested by your doctor to help assess your symptoms and make the diagnosis.

If you are MIBG-positive it means that your tumour is receptive to this chemical - in other words it has special receptors on its surface which take up the MIBG.

MIBG can also be used as a treatment, and is one of the so-called radionuclide therapies.

MIBG combined with a much higher dose of radioactivity than that used for the scan is injected into a vein in your arm via a cannula (a thin plastic tube). It takes approximately 40 minutes to complete the very slow injection (also known as an infusion). The radioactive chemical sticks to the MIBG receptors on the tumours cells and works to kill them off while causing no harm to the healthy cells.

Instead of MIBG therapy occasionally some patients have been treated with other radionuclide therapies. This has only really occurred in a few cases and is not currently standard practice.

Chemotherapy

There is a role for chemotherapy in some patients with malignant tumours that are not suitable for radionuclide therapy or in whom the tumour is growing more quickly and so chemotherapy may work better. The chemotherapy medications used can have side effects and this will be discussed in detail with you prior to considering this therapy.



Inherited Phaeochromocytomas

In about 1 in 4 phaeo patients, the tumour may be associated with a genetic syndrome including multiple endocrine neoplasia type 2 (MEN2).

Patients with MEN2 have a 50% risk of phaeos and almost all will also develop cancer of the thyroid gland in the neck (medullary thyroid cancer). Rarely tumours may occur in the parathyroid glands that lie close to or within the thyroid gland. Initial treatment for MEN2 is for medullary thyroid cancer and consists of the removal of the thyroid gland and surrounding lymph nodes, usually in early childhood. Subsequent management for the other growths associated with MEN2 consists of the monitoring of hormone levels using blood and urine tests, and scans of the neck and abdominal area, sometimes leading to treatment in the form of surgical removal of the tumour and/or affected gland.

What are Paragangliomas?

Paragangliomas are similar to phaeos but are much less common and develop outside of the adrenal glands. They occur more commonly in children rather than adults. Paragangliomas arise from the extra-adrenal paraganglia, which are cells that form part of the nervous system running from the neck to the pelvis, and therefore, tumours can develop anywhere between these two points.

Treatment for Paragangliomas

The management plan is similar to that of phaeos. If surgery can be done to remove as much of the tumour as possible then this should be performed. As with phaeos, patients are given anti-hypertensive medications prior to surgery to reduce the risks of changes in blood pressure during the procedure.

If surgery is not an option or there is some left over tumour that surgeons are not able to remove then these patients can also be considered for treatment with MIBG therapy. It is again dependent on how well the MIBG is taken up by the tumour cells.

Inherited Paraganglioma Syndromes


It is only relatively recently that four types of hereditary paraganglioma syndromes have been identified caused by gene changes in the succinate dehydrogenase (SDH) enzyme. Each child of an affected parent has a 1 in 2 or 50% risk of inheriting the gene change:

SDHD and SDHAF2: these gene changes can only be inherited from an affected father and typically lead to the development of paragangliomas in the head or neck region. Although these tumours usually do not produce hormones, large tumours may cause symptoms such as coughing, hearing loss in one ear, or difficulty swallowing.

SDHC and SDHB: these gene changes can be inherited from either an affected mother or father:

SDHC patients typically develop paragangliomas in the head or neck region that do not usually produce hormones, although large tumours may cause symptoms such as coughing, hearing loss in one ear, or difficulty swallowing.

SDHB patients usually develop extra-adrenal paragangliomas in the abdomen and have a higher risk of their tumours being cancerous and spreading (metastasizing). Like phaeos, these tumours usually secrete hormones called catecholamines that may cause symptoms such as high blood pressure, rapid heartbeat, headaches or sweating. Less commonly these patients may also develop paragangliomas in the head and neck region, as well as kidney and thyroid cancer (papillary thyroid cancer).



In addition, all patients with SDH gene changes are at risk of developing adrenal gland phaeochromocytomas.

Screening for patients with a known SDH gene change or those at 50% risk of inheriting the mutation will undergo similar screening to phaeo patients, including 24 urine collections from around the age of 5 years, annual abdominal ultrasound or MRI scans from around 7 years and neck/thorax MRI every 3 years from age 15. The first treatment of choice is usually surgery.

Medications


To control blood pressure drugs called alpha blockers are commonly used, however, other blood pressure medications may also be required.

If both adrenal glands are removed or you become deficient in the hormones released from the adrenal glands, you will require life-long treatment with replacement steroids (corticosteroids). The two main drugs that a patient must take are hydrocortisone and fludrocortisone. They replace the cortisol and another hormone called aldosterone, which are normally produced by the adrenal glands. The drugs take over in the maintenance of normal blood sugar levels, the promotion of recovery from injury and stress, and the regulation of the balance of mineral salts and water content of the body.

Useful Organisations



NET Patient Foundation
Patient support and advice
0800 434 6476
www.netpatientfoundation.org



AMEND (Association for Multiple Endocrine Neoplasia Disorders)

Support and information for people affected by Multiple Endocrine Neoplasia Disorders and associated endocrine tumours and syndromes
01892 516076
www.amend.org.uk

Further support



NET Patient Foundation

From diagnosis, throughout treatment and beyond, our services are here every step of the way. Here is an overview of all the services we offer to people living with and beyond NET cancer.

Helpline - 0800 434 6476

Our free, confidential helpline is here for anyone who has questions about NET cancers (neuroendocrine tumours). Your call will be answered by one of our nurses or trained staff members with experience of NET cancer. Whatever your concern, you can be confident we will understand the issues you might be facing, and that the information you receive is clear and up-to-date. We will also let you know where else you can go for further support. The helpline is open 10am-2pm Monday to Friday. We also operate a call-back service for those who wish to leave a message out of hours.

Website - www.netpatientfoundation.org

We know how important it is to understand as much as possible about your NET cancer. Our website is here round-the-clock giving you instant access to information when you need it. As well as clinical information, you'll find real life experiences and access to the largest online NET cancer community in the UK, so you can share your questions or concerns with other people in a similar situation.

Discussion forums

Through our discussion forums you can exchange tips on coping with the side effects of treatment, ask questions, share experiences and talk through concerns online. Our dedicated areas for popular topics should make it easy for you to find the information you're looking for. The discussion forums are easy-to-use. If you're feeling anxious or just need to hear from someone else who's been there, they offer a way to gain support and reassurance from others in a similar situation to you.

Ask the nurse

If you find it difficult to talk about your cancer, we can answer your questions by email instead. Our 'ask the nurse' service is available on the website - complete a short form that includes your question and we'll get back to you with a confidential, personal response.

Information and support sessions

We run information and support sessions for people living with NET cancer. These meetings include talks from some of the country's top NET specialists, invaluable Question & Answer sessions, as well as an opportunity to meet other NET patients. For information about meetings in your area, please see our website or call our helpline.

Information resources

Our free information resources are for anyone affected by NET cancer. They are here to answer your questions, help you make informed decisions and ensure you know what to expect. All of our information is written and reviewed regularly by healthcare professionals and people affected by NET cancer, so you can trust the information is up-to-date, clear and accurate. You can order our publications by sending us an email or calling the helpline. All our publications can also be downloaded from our website.

Other organisations

Macmillan Cancer Support

89 Albert Embankment London SE1 7UQ

General enquiries: 020 7840 7840

Helpline: 0808 808 0000

Website: www.macmillan.org.uk

Textphone: 0808 808 0121 or Text Relay

Macmillan Cancer Support provides practical, medical, emotional and financial support to people living with cancer and their carers and families. Over the phone, its cancer support specialists can answer questions about cancer types and treatments, provide practical and financial support to help people live with cancer, and are there if someone just wants to talk. Its website features expert, high-quality information on cancer types and treatments, emotional, financial and practical help, and an online community where people can share information and support. Macmillan also funds expert health and social care professionals such as nurses, doctors and benefits advisers.

The NET Patient Foundation incorporating Living with Carcinoid was formed at the start of 2006 and has Charity Commission status.

The Foundation has the following aims:

- ✦ To provide accurate and up-to-date information for people living with, or affected by, NET cancers
- ✦ To provide support for patients and others affected by NET cancers
- ✦ To provide education for healthcare professionals in the treatment and care of NET patients
- ✦ To raise awareness of NET cancers within the medical community, and amongst the general public
- ✦ To improve access to care for NET patients
- ✦ To raise funds, which will help to support research around the UK & Ireland

Did you know?

- 50% of patients are diagnosed when the cancer has already spread
- Early diagnosis significantly improves outcomes for patients
- Help us to make a difference. Donate now at www.netpatientfoundation.org

Why the moth?

We have chosen the moth as our logo to symbolise this 'camouflaged' condition. We aim to encourage medical professionals to consider an uncommon alternative when symptoms persist (i.e. a NET rather than IBS).

"When you see an insect beating its brightly patterned wings against your window, don't assume it's a butterfly. It could be a moth."

Written by:

Jo Grey AMEND

Professor Ashley Grossman BA, BSc, MD, FRCP, FMedSci

Dr. Raj Srirajaskanthan MD, MRCP



NET Patient Foundation

incorporating Living with Carcinoid

Registered Charity No. 1092386

The NET Patient Foundation supports people diagnosed with
neuroendocrine tumours and their families.

For further information and to make contact telephone 0800 434 6476

or visit our website:

www.netpatientfoundation.org

Published February 2012

delta design and print ltd